

Structure of the X-ray Emission from the Jet of 3C 273

H.L. Marshall¹, D.E. Harris², J.P. Grimes², J.J. Drake², A. Fruscione², M. Juda², R.P. Kraft², S. Mathur³, S.S. Murray², P.M. Ogle¹, D.O. Pease², D.A. Schwartz², A.L. Siemiginowska², S.D. Vrtilek², B.J. Wargelin²
hermanm@space.mit.edu

ABSTRACT

We present images from five observations of the quasar 3C 273 with the Chandra X-ray Observatory. The jet has at least four distinct features which are not resolved in previous observations. The first knot in the jet (A1) is very bright in X-rays. Its X-ray spectrum is well fitted with a power law with $\alpha = 0.60 \pm 0.05$ (where $S_\nu \propto \nu^{-\alpha}$). Combining this measurement with lower frequency data shows that a pure synchrotron model can fit the spectrum of this knot from 1.647 GHz to 5 keV (over nine decades in energy) with $\alpha = 0.76 \pm 0.02$, similar to the X-ray spectral slope. Thus, we place a lower limit on the total power radiated by this knot of 1.5×10^{43} erg/s; substantially more power may be emitted in the hard X-ray and γ -ray bands.

Knot A2 is also detected and is somewhat blended with knot B1. Synchrotron emission may also explain the X-ray emission but a spectral bend is required near the optical band. For knots A1 and B1, the X-ray flux dominates the emitted energy. For the remaining optical knots (C through H), localized X-ray enhancements that might correspond to the optical features are not clearly resolved. The position angle of the jet ridge line follows the optical shape with distinct, aperiodic excursions of $\pm 1^\circ$ from a median value of -138.0° . Finally, we find X-ray emission from the “inner jet” between 5 and $10''$ from the core.

Subject headings: Quasars: individual (3C 273) – galaxies: jets – X-Rays: Galaxies

1. Introduction

Previous high resolution observations of the 3C 273 jet using Merlin at 1.647 GHz and the Hubble Space Telescope (HST) Wide Field Planetary Camera 2 (Bahcall et al. 1995) showed that the overall shape of the jet is somewhat different between the optical and radio bands. The optical image appears dominated by elongated knots roughly $0.1''$ by $0.5''$ in size, while the radio image gives an indication of a “cocoon” structure, especially at the end, or “head”. Bahcall et al. speculated that the cocoon is slowly moving material enveloping a relativistically moving flow. Although relativistic motion is required to explain superluminal motion in the quasar core, it is not yet clear that the flow is relativistic in the jet on a kpc scale.

Harris & Stern (1987) used the *Einstein* Observatory data on 3C 273 to detect X-ray emission from the jet, which was less than 1% of the flux of the core. More recently, Röser et al. (2000), examined ROSAT HRI images at $\sim 5''$ resolution to show that the X-ray emission drops with distance along the jet. Using models of the profile along the jet and multi-color ground-based images at $1.3''$ resolution, they generated spectral energy distributions (SEDs) for knots in the jet and found that the X-ray flux of knot A1⁴ is consistent with that expected by extrapolating a simple synchrotron model from the radio through the optical with one population of electrons. The highest energy electrons in their model had $\gamma > 10^7$. Synchrotron self-Compton (SSC) calculations generally fail to predict X-ray intensities commensurate with those observed for any of the knots.

¹Center for Space Research, Massachusetts Institute of Technology

²Harvard-Smithsonian Center for Astrophysics

³Ohio State University

⁴We will refer to the knot naming convention given by Bahcall et al. (1995) (reproduced in Fig. 1).

X-ray images with the Chandra X-ray Observatory are now resolving the spatial structure along quasar jets. The first such image, of the quasar PKS 0637–752, proved remarkable because of the strong X-ray fluxes of the jet knots, relative to the radio fluxes (Schwartz et al. 2000). Simple synchrotron and thermal models could be ruled out easily while SSC models required unreasonable conditions. Tavecchio et al. (2000) and Celotti, et al. (2000) have suggested that inverse Compton scattering of the cosmic microwave background could produce the required X-ray fluxes and require the jet material to be moving relativistically at a small angle to the line of sight as in models of the core. We present Chandra images and spectra for the jet in 3C 273 that we examine in light of these models.

2. Observations and Analysis

2.1. X-ray Data Reduction

3C 273 was observed three times as part of the calibration of the Chandra grating spectrometers, once in a direct imaging mode as part of science verification, and twice more as the gratings failed to insert during calibration observations. See Table 1 for a list of observations, totalling 193,230 s of exposure time. Results from the dispersed spectra of the core are still being analyzed as part of the ongoing effort to verify the grating spectrometer effective area calibration and will be presented elsewhere. The Low Energy Transmission Grating (LETG) and High Resolution Camera (HRC) combination gave significantly fewer counts in the jet compared to the others, so was not used in the combined X-ray image. We present here the results from the zeroth order portion of the ACIS grating observations and combine these data with that of the HRC-I and ACIS-S imaging observations for image analysis. The dispersed spectra of the jet were extremely faint; we determined that there were no significant emission lines and did not examine these data any further.

Each of the grating observations had well-known artifacts that distort the zeroth order images. Two artifacts are relevant to observations of the jet. Bright sources observed with ACIS will show up with a “readout streak” (see the Chandra Proposers’ Observatory Guide, or POG). In no ACIS observation, however, did the streak affect the image of the jet. The LETG has two support structures that produce diffraction patterns (see the POG). This pattern is spaced at 60° intervals and, again, did not interfere with the im-

age of the jet. Individual observations have slightly uncertain absolute pointing, so we repositioned each image separately. The HRC and LETG/ACIS observations were combined by referencing to the core. Due to pileup, the two non-grating ACIS observations and the High Energy Transmission Grating observations had zeroth order images that were severely affected by pileup so the first knot was used as a reference. We estimate the uncertainty in this procedure gives a relative offset between the core and the jet of less than $0.05''$ by examining the location of the center of the wings of the core and comparing to the position of the first knot in each data set.

Figure 1 shows the combined X-ray image binned in $0.2''$ pixels. The jet shows clear curvature but mostly has a one-dimensional appearance, so a profile was computed for quantitative analysis. The X-ray profile (Fig. 2) was derived by summing data in a $1.5''$ wide window centered on a position angle of -138.0° , which is the centerline of the jet. We estimate that the correction for aperture losses is $10 \pm 5\%$ at any location along the jet, based on using a wider extraction region. The peak of emission occurs just under $13''$ from the core. Fitting a Gaussian profile to it, the centroid is $12.''93 \pm 0.''01$ from the core and the dispersion of a Gaussian is $0.''33 \pm 0.''01$ (for a FWHM of $0.78''$). The 2σ limit to the FWHM of this knot is $0.''3$, given that point sources have a projected FWHM of $0.''75$, which is determined from readout streak data for the core of 3C 273 and other bright point sources. The next peak in the X-ray emission is clearly extended, from 14.0 to $15''$ from the core, dropping more steeply on the downstream side. From $16''$ to $21''$ the X-ray emission appears somewhat devoid of distinct features but with some possible surface brightness variations; there may be an unresolved knot at $20''$. The X-ray flux reaches the background level at $21''$ from the nucleus.

Several regions were selected for X-ray spectral fitting: a) a $1''$ radius circle centered on the first bright knot, b) a similar circle centered on the extended knot at $15''$ and c) a rectangular box extending from $16''$ to the end of the jet. The 0.5–8 keV ACIS-S spectra were combined and fitted with power law models holding N_H fixed at $1.71 \times 10^{20} \text{ cm}^{-2}$ (Albert et al. 1993). The resultant spectral indices were $\alpha_a = 0.60 \pm 0.05$, $\alpha_b = 0.88 \pm 0.07$, and $\alpha_c = 0.75 \pm 0.05$ (where $S_\nu \propto \nu^{-\alpha}$); no spectral evolution is detected along the jet. Flux densities of several knot regions are given in Table 2. We estimate that the total jet power is about 0.4% of the core power in the 0.5–5.0 keV band.

X-ray emission is just detectable between the core and the knot A1, as shown in Fig. 3. The 5-10'' annulus shows a peak at the position angle of the 10-20'' jet in addition to peaks at the position angles of the readout streak. It is difficult to quantify precisely the flux of the inner jet due to the ripple inherent in azimuthal profiles this close to a bright source which is caused by mirror support structure. Accounting for the ripple by fitting a sinusoid to the local background, we estimate that the count rate from the inner jet is about 0.012 ± 0.001 count s⁻¹, corresponding to a total flux density of 6.9 ± 0.6 nJy at 1 keV for a power law spectrum with $\alpha = 0.6$.

2.2. Comparison to the Optical Emission

X-ray components were identified using images obtained from the HST archives. Fluxes were determined in filters F450W, F622W, F814W, and F160W (NICMOS).⁵ The planetary camera (PC) observation using filter F622W was used for direct comparisons to the X-ray jet profile, so the centroid of the quasar was determined by isolating the image diffraction spikes (Bahcall et al. 1995), fitting these with lines, and determining the intersection of the two lines (only two spikes were available from this image). Spatial distortions were corrected using polynomial coefficients given by Holtzman et al. (1995). The optical profile was obtained using the same method as used for the X-ray profile and is also shown in Fig. 2. An offset of 0.22'' was found between the centroids of the the first X-ray peak and the peak of knot A1 from the PC image. The positional uncertainty is dominated by systematic uncertainties in measurement of the quasar core, so this difference between the X-ray and optical positions of this bright knot are not likely to be significant. The optical emission of A1 is clearly extended along a position angle closely aligned to the overall PA of the jet (see Fig. 1); the profile is well fitted by a Gaussian with $\sigma = 0.3''$ which is consistent with the X-ray profile at the 2σ level.

Identifying the sources of the remaining X-ray emission is not quite so straightforward as for knot A1. Region b (the extended knot at 15'' from the core) is not consistent with a single point source at knot B1 but is likely to be a blend of point sources at knots B1 and A2, a somewhat weaker knot in the HST image.

There appears to be a discrete source of X-ray emission near the positions of knots D and H3. More data are needed to tell if knots besides these – i.e., C1, C2, and C3 – are also discrete sources of X-ray emission.

3. Discussion

3.1. Morphology

The overall shape of the jet is quite similar in the optical and X-ray bands: there is distinct curvature and the lengths are about the same. The optical emission between the knots is $\gtrsim 0.5''$ wide (Bahcall et al. 1995) and the X-ray emission is marginally consistent with this level of broadening. There are several important differences, however. Knot A1 is much more prominent in the X-ray data than in the optical image. Furthermore, the X-ray jet fades along the jet while the optical knots have similar brightnesses. Röser et al. (2000) also noted this difference.

The X-ray emission from knot A1 is consistent with a point source but we cannot yet exclude the possibility that it is as extended as the optical emission along the jet axis. The Merlin map shows that this knot is similarly extended in the 1.647 GHz band. The radio emission of the other knots is quite difficult to discern within the radio cocoon, giving rise to the impression that bulk of the radio emission is physically distinct from the optical and X-ray emission regions. We find no significant X-ray emission from either extensions (inner or outer, see Fig. 1), lending support to the interpretation that these are unrelated to the jet (Röser & Meisenheimer 1991).

3.2. Spectra

The overall spectral energy distribution (SED) for knot A1 (Fig. 5) appears to fit a simple synchrotron model, as suggested by Röser et al. (2000). Our estimate for the X-ray flux density is $\sim 2\times$ higher, suggesting that the uncertainties derived from the ROSAT data were underestimated. Amazingly, the flux of this knot fits the overall slope of the SED (based on table 2), 0.76 ± 0.02 , to within the uncertainties, even after extrapolating over several orders of magnitude in frequency. Furthermore, the spectral slope in the X-ray band is similar to that of the SED. Thus, the spectrum does not appear to break within the Chandra bandpass, which is consistent with the excellent spectral fit to the Chandra data. The luminosity of knot A1, 1.5×10^{43} erg/s (for $q_0 = 0.5$ and $H_0 = 70$ km/s/Mpc),

⁵Based on observations made with the NASA/ESA Hubble Space Telescope, obtained from the data archive at the Space Telescope Science Institute. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555.

is about 40% of the total X-ray emission from jet, so if the spectrum of knot A1 extends out to 100-200 keV with $\alpha = 0.6$, then its X-ray flux will dominate the total power of the jet. If the synchrotron break is above 5 keV then $\gamma > 4 \times 10^7$ for the electrons, if one synchrotron model is to fit all the data. The magnetic field is $80\mu\text{G}$, based on minimum energy arguments for nonrelativistic bulk motion. For a cylindrical emitting volume of the size defined by the optical emission, the SSC emission from knot A1 would be less than 0.1 nJy, well below the observed value.

Connecting the radio and optical/X-ray bands of the B1 SED requires a slight bend, consistent with the apparent flattening in the optical band. While the form of the electron energy distribution may not be a pure power law, no spectral cutoff is observed in the SED, so the electron energies may well reach energies comparable to those in knot A1. Röser et al. determined that SSC models of knot B1 would not give rise to such large X-ray fluxes and we confirm this conclusion. As in knot A1, νS_ν is much higher in the X-ray band than in the optical and radio bands. Because $\alpha < 1$ in the X-ray band, we do not yet know where the total powers of these knots peak.

Without detecting spectral cutoffs as observed in the first knot of the jet in PKS 0637–752 (Schwartz et al. 2000), we cannot tell if there is a problem with the synchrotron model for knots A1 and B1 as found in that source. Similarly, the X-ray emission mechanism in the inner jet region is difficult to model without spectra from the radio and optical bands. Although there is no specific evidence that relativistic motion is required to explain the X-ray fluxes of the inner jet or knots A1 and B1, as in the model suggested by Tavecchio et al. (2000) and Celotti, et al. (2000), this beaming model is a promising explanation for the X-ray fluxes of the weaker knots so it could also provide an alternative to the synchrotron models for knots A1 and B1. Thus, based on morphological similarities between the X-ray and optical images and considering that much of the jet power may be dissipated in the knots, we speculate that the knots are locations of internal shocks in a relativistic jet flow that is decelerating before equilibrating with the ambient medium. Alternatively, a helical jet structure would have regularly spaced inflection points where the local jet flow is close to the line of sight; beaming would be enhanced, giving rise to the observed knots.

We thank Herman-Josef Röser for discussions

about the HST data for which he was the principal investigator. We thank Tom Muxlow of Jodrell Bank for providing the Merlin image. This research is funded in part by NASA contracts NAS8-38249, NAS8-39073, and SAO SV1-61010.

REFERENCES

- Albert, C.E., Blades, J. C., Morton, D. C. Lockman, Felix J., Proulx, M. Ferrarese, L. 1993, ApJS, 88, 81
- Bahcall et al. 1995, ApJ, 452, L91
- Celotti, A., Ghisellini, G., and Chiaberge, M. 2000, MNRAS, submitted, (astro-ph/0008021)
- Harris, D.E. & Stern, C.P. 1987, ApJ, 313, 136
- Holtzman et al. 1995, PASP, 107, 156
- Röser, H.-J., and Meisenheimer, K. 1991, A&A, 252, 458
- Röser, H.-J., Meisenheimer, K., Neumann, M., Conway, R.G., and Perley, R.A. 2000, A&A, 360, 99
- Schwartz et al. 2000, ApJ, 540, L69
- Tavecchio, F., Maraschi, L., Sambruna, R., and Urry, C.M. 2000, ApJ, submitted, (astro-ph/0007441)

See color image: fig1.ps.

Fig. 1.— Images of the jet in 3C 273 in three different bands. *Left*: Image at 1.647 GHz using the Merlin array, kindly provided by Tom Muxlow of Jodrell Bank. *Middle*: The Hubble Space Telescope Planetary Camera image in the F622W filter (centered at 6170 Å). Features are labelled according to the nomenclature used by Bahcall et al. (1995). *Right*: Raw Chandra image of the X-ray emission from the jet of 3C 273 in 0.1'' bins overlaid with a version of the HST image smoothed with a Gaussian profile in order to match the X-ray imaging resolution. The X-ray and optical images have been registered to each other to about 0.05'' using the position of knot A1. The overall shape of the jet is remarkably similar in length and curvature but the X-ray emission fades to the end of the jet so individual C knots are not discernable. Other differences are more apparent in Fig. 2. The radio emission is much fainter at knot A1 and is displayed with a logarithmic scaling.




Fig. 2.— Profiles of the 3C 273 jet in the X-ray (histogram, in counts per $0.1''$ bin) and optical bands. The optical data are from a Planetary Camera image in filter F622W, taken from the Hubble Space Telescope (HST) archive, are scaled to match the X-ray histogram for knot A1: $0.325 \mu\text{Jy}$ per $0.04554''$ bin at a vertical value of 500. The raw optical profile was smoothed with a Gaussian with $\sigma = 0.25''$ (FWHM = $0.60''$) and was scaled to $0.27 \mu\text{Jy}$ per $0.04554''$ bin at a vertical value of 500. The optical profiles were displaced $0.22''$ closer to the core to provide a better match between the X-ray and optical profiles of knot A1; systematic registration uncertainties are of this order. Beyond knots A1 and B1 ($12.9''$ and $15.0''$ from the core, respectively), other knots are not clearly detected individually in the X-ray profile.




Fig. 3.— Azimuthal histograms from two different annuli centered on the quasar core. The top panel, for the annulus from 12-15'' from the core, shows the jet at a position angle of -138° (shown by dashed lines) and the readout streaks at about 110° and -70° (shown by dotted lines). The bottom panel shows the histogram for the 5-10'' annulus. The two brightest peaks are at the position angles of the readout streak while the third brightest is at the same position angle as the large scale jet. The ripple is due the mirror support structure, which causes shadows every 30° . The peak at -138° is a detection of X-rays from the “inner jet”, which is not very bright optically but is detected in the radio band. The position angle of the inner jet is rotated slightly to the north compared to the 5-10'' portion of the jet. A similar rotation is observed in the Merlin map (see figure 1).




Fig. 4.— Two ACIS-S spectra that were simultaneously fit with a power law model ($\alpha = 0.60$; solid lines) including Galactic absorption of $N_H = 1.71 \times 10^{20}$ atoms/cm². Only data within 0.5-8 keV were included in the analysis. The fitting was performed on the original 1024 PI bins using Cash statistics in Sherpa. We rebinned the data to have minimum of 10 counts per bin for the plotting purposes only. The upper data set, from observation ID 1711, is displayed in the original scale, while the lower data set, from observation ID 1712, was rescaled downward by $\times 10$ for clarity.




Fig. 5.— The spectral energy distributions (SEDs) of the 3C 273 jet knots A1, B1/B2, D/H3, and the inner jet. The overall spectrum of knot A1 fits a simple power law, shown as the dashed line with a slope of 0.25 so the spectrum is consistent with a simple synchrotron model with a single power law distribution of electrons. For knot B1, however, the SED appears to flatten between the radio and optical bands; νS_ν still peaks in the X-ray band. A single synchrotron model does not fit the SED of the D/H3 region.

Table 1: Chandra Observations of 3C 273

Obs. ID	Detector	Grating	Date	Exposure (s)
459	ACIS-S	HETG	10 Jan 2000	38600
460	HRC-I	none	22 Jan 2000	20260
461	HRC-S	LETG	9 Jan 2000	40280
1198	ACIS-S	LETG	9–10 Jan 2000	38160
1711	ACIS-S	none	14 Jun 2000	28130
1712	ACIS-I	none	14 Jun 2000	27800

Table 2: Fluxes of Knots in the 3C 273 Jet

Frequency (Hz)	Flux Density (μ Jy)		
	A1	B1	D/H3
1.65×10^9	$4.2 \pm 1.2 \times 10^4$	$4.5 \pm 2.0 \times 10^5$	$6.5 \pm 1.0 \times 10^5$
1.87×10^{14}	10.7 ± 2.3	14.8 ± 8.1	71.3 ± 10.7
3.76×10^{14}	6.7 ± 2.0	6.9 ± 1.8	13.2 ± 2.8
4.85×10^{14}	5.15 ± 0.83	5.16 ± 0.97	8.2 ± 1.2
6.59×10^{14}	4.32 ± 0.91	4.11 ± 0.90	4.58 ± 1.02
2.42×10^{17}	0.038 ± 0.004	0.023 ± 0.002	0.0083 ± 0.0008

